



(11) Publication number : **0 518 818 A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number : **92810422.3**

(51) Int. Cl.⁵ : **C07C 257/18, A61K 31/155**

(22) Date of filing : **02.06.92**

(30) Priority : **11.06.91 US 714109**

(43) Date of publication of application :
16.12.92 Bulletin 92/51


(84) Designated Contracting States :
AT BE CH DE DK ES FR GB GR IT LI LU NL PT SE

(71) Applicant : **CIBA-GEIGY AG**
Klybeckstrasse 141
CH-4002 Basel (CH)

(72) Inventor : **Main, Alan Joseph**
60 Briar Lane
Basking Ridge, NJ 07920 (US)

(54) Arylethers, their manufacture and use as medicament.

(57) The invention relates to the compounds of the formula



and pharmaceutically acceptable salts thereof, in which :

R is hydrogen or an acyl radical which is derived from an organic carbonic acid, an organic carboxylic acid, a sulfonic acid, or a carbamic acid ;

R₁ is a substituent selected from an aliphatic hydrocarbon radical, an araliphatic hydrocarbon radical and a cycloaliphatic hydrocarbon radical ;

X₁ and X₃, independently of one another, are oxygen (-O-) or sulphur (-S-) ; and

X₂ is a divalent aliphatic hydrocarbon radical which may be interrupted by an aromatic radical ;

wherein the phenyl rings of formula I may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid ;


wherein aryl in the above definitions may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid ; and wherein a cycloaliphatic hydrocarbon radical may be substituted by an aliphatic radical. The compounds are useful as selective LTB₄ receptor antagonists in the treatment of conditions or syndromes in mammals which are responsive to LTB₄ receptor antagonism.

The invention relates to the phenoxyalkoxyphenyl derivatives and thio analogs as defined herein which are particularly useful as selective Leukotriene B₄ (LTB₄) receptor antagonists, methods for preparation thereof, pharmaceutical compositions comprising said compounds, and a method of antagonizing LTB₄ and of treating conditions or syndromes in mammals which are responsive to LTB₄ antagonism using said compounds or pharmaceutical compositions comprising said compounds of the invention.

Leukotriene B₄ (LTB₄) is an important inflammatory mediator being a potent chemotactic agent and activator of polymorphonuclear leucocytes (PMN's) and monocytes. It modulates the production and effects of other important inflammatory mediators e.g. Interleukin-1 and gamma Interferon. LTB₄ has been implicated in the pathogenesis of a number of inflammatory diseases, such as rheumatoid arthritis, inflammatory bowel disease, psoriasis, non-steroidal-antiinflammatory-drug-induced gastropathy, adult respiratory distress syndrome (ARDS), myocardial infarction, allergic rhinitis, hemodialysis-induced neutropenia, and late phase asthma.

There is a strong need in the art in finding potent antagonists of LTB₄ on human PMN's, especially those which are orally active. It has been found that the compounds according to the present invention exhibit significant LTB₄ antagonistic activity on human PMN's and are orally active.

The invention relates to phenoxyalkoxyphenyl derivatives and thio analogs of the formula



and pharmaceutically acceptable salts thereof, in which:

R is hydrogen or an acyl radical which is derived from an organic carbonic acid, an organic carboxylic acid, a sulfonic acid, or a carbamic acid;

R₁ is a substituent selected from an aliphatic hydrocarbon radical, an araliphatic hydrocarbon radical and a cycloaliphatic hydrocarbon radical;

X₁ and X₃, independently of one another, are oxygen (-O-) or sulphur (-S-); and

X₂ is a divalent aliphatic hydrocarbon radical which may be interrupted by an aromatic radical;

wherein the phenyl rings of formula I may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid;

wherein aryl in the above definitions may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid;

wherein a cycloaliphatic hydrocarbon radical may be substituted by an aliphatic radical;

which are particularly useful as selective LTB₄ antagonists, methods for preparation thereof, pharmaceutical compositions comprising said compounds, and a method of antagonizing LTB₄ and of treating diseases in mammals which are responsive to LTB₄ antagonism using said compounds or pharmaceutical compositions comprising said compounds of the invention.

As the compounds according to the invention have a basic centre, they can thus form acid addition salts, especially pharmaceutically acceptable salts. These are formed, for example, with inorganic acids, such as mineral acids, for example sulfuric acid, a phosphoric or hydrohalic acid, or with organic carboxylic acids, such as (C₁-C₄)-alkanecarboxylic acids which, for example, are unsubstituted or substituted by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, for example aspartic or glutamic acid, benzoic acid or with organic sulfonic acids, such as (C₁-C₄)-alkane- or arylsulfonic acids which are unsubstituted or substituted, for example, by halogen, for example methane- or toluenesulfonic acid. Preferred are salts formed with hydrochloric acid, methanesulfonic acid and maleic acid.

An acyl radical which is derived from an organic carbonic acid is, for example, alkoxycarbonyl or alkenyloxycarbonyl which in each case are unsubstituted or substituted by an aromatic radical or is cycloalkoxycarbonyl which unsubstituted or substituted by lower alkyl.

An acyl radical which is derived from an organic carboxylic acid is, for example, lower alkanoyl, phenyl-lower alkanoyl or unsubstituted or substituted aroyl, such as benzoyl, naphthoyl, indanoyl or fluorenyl.

An acyl radical which is derived from a sulfonic acid is, for example, alkanesulfonyl, arylalkanesulfonyl, cycloalkanesulfonyl or arylsulfonyl.

5 An acyl radical which is derived from a carbamic acid is, for example, amino-carbonyl which is substituted by alkyl, arylalkyl or aryl.

An aromatic radical is, for example, unsubstituted or substituted such as monosubstituted or polysubstituted, for example, disubstituted or secondarily trisubstituted, carbocyclic aryl, such as phenyl, naphthyl, indanyl or fluorenyl.

10 An aliphatic hydrocarbon radical is, for example, lower alkyl, lower alkenyl and secondarily lower alkynyl.

An araliphatic hydrocarbon radical is, for example, optionally substituted phenyl-lower alkyl and secondarily phenyl-lower alkenyl and phenyl-lower alkynyl.

A cycloaliphatic hydrocarbon radical is, for example, cycloalkyl and secondarily cycloalkenyl, which is unsubstituted or mono- or polysubstituted, for example, disubstituted, by lower alkyl. A cycloaliphatic hydrocarbon radical is also optionally substituted cycloalkyl-lower alkyl.

15 A divalent aliphatic hydrocarbon radical is, for example, lower alkylene.

A divalent aliphatic hydrocarbon radical which is interrupted by an aromatic radical is, for example, lower alkylene-phenylene-lower alkylene or lower alkylene-naphthylene-lower alkylene.

20 An aliphatic alcohol is, for example, a lower alkanol or lower alkenol, and an araliphatic alcohol is, for example, a phenyl-lower alkanol.

Hydroxy which is etherified by an aliphatic, cycloaliphatic or araliphatic alcohol is, for example, lower alkoxy or lower alkenyloxy, cyclohexyl-lower alkoxy, and phenyl-lower alkoxy.

An aliphatic carboxylic acid is, for example, a lower alkanic or lower alkenic acid, and an araliphatic carboxylic acid is, for example, a phenyl-lower alkanic acid.

25 Hydroxy which is esterified by an aliphatic or araliphatic carboxylic acid is, for example, lower alkanoyloxy, secondarily lower alkenoyloxy, or is phenyl-lower alkanoyloxy.

The phenyl rings of formulae I and IA as well as aromatic radicals referred to before and hereafter are generally unsubstituted or further substituted such as monosubstituted or polysubstituted, for example disubstituted or secondarily trisubstituted, in particular, for example, by a substituent selected from the group consisting of halogen, trifluoromethyl, lower alkyl, lower alkenyl, lower alkynyl, hydroxy, lower alkoxy, lower alkenyloxy, cyclohexyl-lower alkoxy, phenyl-lower alkoxy, lower alkanoyloxy, lower alkenoyloxy, and phenyl-lower alkanoyloxy. Preferably, the phenyl rings of formula I and IA do not exhibit any additional substituent.

Preferred positions of the following structural elements in the corresponding phenyl ring in formula I are: position 2 (ortho) for $-OR_1$ of the corresponding phenyl ring with respect to

35 $-X_1-$, position 4 (para) for $-C(=NH)-NH-R$ of the corresponding phenyl ring with respect to $-X_3-$.

The substituent R may be located on either nitrogen of the $-C(=NH)NH_2$ grouping.

The general definitions used above and below have, if not defined differently, the following meanings:

The expression "lower" means that corresponding groups and compounds in each case contain in particular not more than 7, preferably not more than 4, carbon atoms.

40 Alkoxy-carbonyl is, in particular, C_2-C_{12} -alkoxy-carbonyl and is, for example, methoxy-, ethoxy-, propyloxy-, pivaloyloxy- or octyloxy-carbonyl. C_2-C_8 -alkoxy-carbonyl is preferred.

Alkenyloxy-carbonyl is, in particular, C_3-C_{12} -alkenyloxy-carbonyl, for example, allyloxy-carbonyl. Preferred is C_3-C_6 -alkenyloxy-carbonyl.

45 Cycloalkyloxy-carbonyl is, in particular, C_3-C_7 -cycloalkoxy-carbonyl, preferred is cyclopentyloxy-carbonyl or cyclohexyloxy-carbonyl.

Lower alkanoyl is, in particular, C_1-C_7 -alkanoyl and is, for example, formyl, acetyl, propionyl, butyryl, isobutyryl or pivaloyl. C_2-C_5 -Alkanoyl is preferred.

Phenyl-lower alkanoyl is, in particular, phenyl- C_2-C_7 -alkanoyl and is, for example, phenylacetyl or 2- or 3-phenylpropionyl. Phenyl- C_2-C_4 -alkanoyl is preferred.

50 Naphthoyl is 1- or 2-naphthoyl.

Indanoyl is, for example, 1-, 2-, 3- or 4-indanoyl.

Fluorenyl is, for example, 1-, 2-, 3-, 4- or 5-fluorenyl.

Alkanesulfonyl is, in particular, C_1-C_7 -alkanesulfonyl and is, for example, methane-, ethane-, n-propane- or isopropanesulfonyl. C_1-C_4 -alkanesulfonyl is preferred.

55 Arylalkanesulfonyl is, in particular, phenyl- C_1-C_7 -alkanesulfonyl, for example, benzyl- or 1- or 2-phenylethan-sulfonyl. Phenyl- C_1-C_4 -alkane-sulfonyl is preferred.

Cycloalkanesulfonyl is, in particular, C_3-C_7 -cycloalkanesulfonyl, preferred is cyclopentanesulfonyl or cyclohexanesulfonyl.

Naphthyl is 1- or 2-naphthyl.

Indanyl is, for example, 1-, 2-, 3- or 4-indanyl.

Fluorenyl is, for example, 1-, 2-, 3-, 4- or 5-fluorenyl.

Lower alkyl is, in particular, C₁-C₇ alkyl and is, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl and furthermore includes corresponding pentyl, hexyl and heptyl radicals. C₁-C₄-alkyl is preferred.

Lower alkenyl is, in particular, C₃-C₇ alkenyl and is, for example, 2-propenyl or 1-, 2- or 3-butenyl. C₃-C₅-alkenyl is preferred.

Lower alkynyl is, in particular, C₃-C₇ alkynyl and is preferably propargyl.

Phenyl-lower alkyl is, in particular, phenyl-C₁-C₄-alkyl and is preferably benzyl, 1- and 2-phenethyl, while phenyl-lower alkenyl and phenyl-lower alkynyl are, in particular, phenyl-C₂-C₆ alkenyl and -alkynyl, in particular 2-phenyl-vinyl, 3-phenylallyl and 3-phenylpropargyl.

Cycloalkyl is, in particular, C₃-C₇ cycloalkyl and is, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. Cyclopentyl and cyclohexyl are preferred.

Cycloalkenyl is, in particular, C₃-C₇ cycloalkenyl and is preferably cyclopent-2- or -3-en-yl, or cyclohex-2- and -3-en-yl.

Lower alkylene e.g. in amino which is disubstituted by lower alkylene is, in particular, C₂-C₆-alkylene and is, for example, butylene, pentylene, or 2,6-butylene. Preferred is C₄-C₆-alkylene, especially pentylene.

Lower alkylene (X₂) is, in particular, C₂-C₈-alkylene, preferably straight-chain, and is, for example, ethylene, propylene, butylene, pentylene, hexylene, and also octylene. C₄-C₇-alkylene is preferred, especially pentylene and also butylene, hexylene or heptylene.

Lower alkylene which is interrupted by a phenyl radical (X₂) is, in particular, lower alkylene-phenylene-lower alkylene or lower alkylene-naphthylene-lower alkylene such as C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene or C₂-C₄-alkylene-naphthylene-C₂-C₄-alkylene, preferably straight-chain, and is, for example, methylene-phenylene-methylene, 1,2-ethylene-phenylene-1,2-ethylene, such as 1,2-ethylene-1,4-phenylene-1,2-ethylene, 1,3-propylene-phenylene-1,3-propylene, such as 1,3-propylene-1,4-phenylene-1,3-propylene, or butylene-phenylene-butylene radicals, also a corresponding 1,2-ethylene-naphthylene-1,2-ethylene radical. C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene or C₂-C₃-alkylene-naphthylene-C₂-C₃-alkylene is preferred, especially 1,2-ethylene-1,4-phenylene-1,2-ethylene.

Halogen is, in particular, fluorine, chlorine or bromine, and furthermore includes iodine.

Lower alkoxy is, in particular, C₁-C₇ alkoxy and is, for example, methoxy, ethoxy, n-propyloxy, isopropyloxy, n-butyloxy, isobutyloxy, sec-butyloxy, tert-butyloxy and furthermore includes corresponding pentyloxy, hexyloxy and heptyloxy radicals. C₁-C₄-alkoxy is preferred.

Lower alkenyloxy is, in particular, C₃-C₇ alkenyloxy and is, for example, allyloxy or but-2-en- or but-3-en-yloxy. C₃-C₅-alkenyloxy is preferred.

Phenyl-lower alkoxy is, in particular, phenyl-C₁-C₄-alkoxy, such as benzyloxy, 1- or 2-phenylethoxy, or 1-, 2- or 3-phenylpropyloxy.

Lower alkanoyloxy is, in particular, C₂-C₈-alkanoyloxy, in particular, C₂-C₆-alkanoyloxy, such as acetyloxy, propionyloxy or pivaloyloxy.

Lower alkenoyloxy is, in particular, C₃-C₈-alkenoyloxy, in particular, C₃-C₅-alkenoyloxy, such as propenoyloxy.

Phenyl-lower alkanoyloxy is, in particular, phenyl-C₂-C₈-alkanoyloxy, in particular, phenyl-C₂-C₆-alkanoyloxy, such as phenylacetyloxy, phenylpropionyloxy or phenylpivaloyloxy.

The compounds of the invention exhibit valuable pharmacological properties in mammals, and are particularly useful as selective Leukotriene B₄ (LTB₄) receptor antagonists, e.g. for the treatment of a condition or syndrome in a mammal responsive to the selective antagonism of LTB₄ receptors, such as rheumatoid arthritis, inflammatory bowel disease, psoriasis, non-steroidal-antiinflammatory-drug-induced gastropathy, adult respiratory distress syndrome (ARDS), myocardial infarction, allergic rhinitis, hemodialysis-induced neutropenia, and late phase asthma. The compounds of the invention are also useful as analgesics for the treatment of pain, of any origin, and for the treatment of osteoarthritis.

The above-cited properties are demonstrable in in vitro and in vivo tests, using advantageously mammals, e.g. rats. Said compounds can be applied in vitro in the form of solutions, e.g. preferably aqueous solutions, and in vivo either enterally or parenterally, advantageously orally, e.g. as a suspension or in aqueous solution. The dosage in vitro may range between about 0.5 ng/ml and about 100 ng/ml. The dosage in vivo may range, depending on the route of administration, between about 1 and about 1000 mg/kg per day.

Beneficial effects are evaluated in pharmacological tests generally known in the art, e.g. as illustrated herein.

Receptor Binding with [³H]-LTB₄ to Intact Human Neutrophils:

Neutrophils (PMN's) are prepared from uncoagulated human venous blood. Blood is dispersed into 50 ml polypropylene tubes containing 15 ml of HESPAN (Dupont, Wilmington, DE), and mixed. Tubes are allowed to stand at room temperature for 40 minutes until most of the red blood cells sediment. The supernatants are removed and centrifuged for 5-10 min at 400 x g. The remaining pellets are diluted in 70 ml of Phosphate Buffered Saline without calcium and magnesium (PBS without metals; GIBCO, Grand Island, NY) and 35 ml of this suspension are placed in each of two polypropylene tubes containing 15 ml of Ficoll-Paque (Sigma, St. Louis, MO). Gradients are then centrifuged for 15 minutes at 420 x g. The mononuclear cell layer is discarded and the remaining red blood cell pellet is resuspended in 10 ml of PBS without metals. Twenty ml of filtered deionized water are added to the suspension for approximately 20 sec followed by the same volume of buffer at two times the normal concentration. The cell suspension is mixed and centrifuged for 5 min at 200 x g, followed by one wash with buffer, and final resuspension.

Binding of [³H]LTB₄ to LTB₄ receptors is measured in intact human polymorphonuclear leukocytes, as described by Gorman and Lin (Gorman, R. and Lin, A Methods Enzymol. 141: 372-378, 1987). Intact human neutrophils are suspended in Hank's Balanced Salt Solution (HBSS) at a concentration of 3 x 10⁶ cells/assay tube. An aliquot of the cell suspension (300 µl) is added to triplicate tubes containing 50 µl [³H]LTB₄ (specific activity 32 Ci/mmol, DuPont-NEN, Boston, MA) at a final concentration of 0.5 nM, 100 µl buffer and 50 µl drug or buffer. Nonspecific binding is determined in the presence of 300 nM LTB₄. The reaction is initiated by addition of cell suspension and continued at 0°C for 20 min. Bound radioactivity is isolated by vacuum filtration through Whatman GF/C glass fiber filters using a Brandel cell harvester and unbound radioactivity removed with 2 x 5 ml washes with ice-cold saline. Filters are placed in polyethylene scintillation mini-vials to which is added 3.5 ml of formula-989 scintillation cocktail (NEN). After equilibration, radioactivity determinations and data calculations are performed using non-linear regression analysis on RS/1.

LTB₄-Induced PMN Aggregation

Human PMNs are prepared as previously described. Neutrophil aggregation is assessed by monitoring the intensity of light passing through a suspension of cells (Craddock et al., J. Clin. Invest. 60: 260-264, 1977) using a Payton dual channel aggregometer (model 300BD). Cuvettes containing 0.25 ml of cell suspension (25 x 10⁶ cells/ml) in PBS without calcium and magnesium are incubated with 5 µg/ml of cytochalasin B for 2 minutes at 37°C. 5 µl of 2 µM LTB₄ in PBS (20 nM final concentration) are added and the aggregation response monitored for 3-5 min, the time required for optimal response. Compounds are solubilized in 0.01M DMSO and then diluted in PBS to 0.001 M. 5 µl of compound solution is added along with cytochalasin B and cells as described above. Following the preincubation period 5 µl of 2 µM LTB₄ are added and aggregation is measured. Percent inhibition of aggregation is calculated by comparing peak heights in the presence and absence of compound. Percent inhibition is plotted as a function of the log concentration of compound and the IC₅₀ determined directly from the graph.

LTB₄-Induced Neutropenia in the Rat

Male Sprague Dawley rats (cr: CDBR; Charles River, Wilmington, MA.) (250-300) grams are fasted overnight prior to the experiment. At least six animals are used per treatment group. Rats are given vehicle or compound either intravenously or orally and at intervals after dosing, neutrophil counts are determined from blood samples obtained just prior to and 20 seconds after intravenous infusion of 200 ng LTB₄. In studies are compound is administered orally, drug is given by gavage. When drug is administered intravenously, rats are first anesthetized with 50 mg/kg i.p. of Sodium Pentobarbital. The jugular vein is exposed and cleaned of the surrounding tissue. At 3, 4 or 18 hours following administration of compound or vehicle by either route, blood samples are taken (0.3 ml of blood in 1.5 ml polypropylene microcentrifuge tube containing 0.01 ml 7.5% EDTA). Blood neutrophil counts are determined using a Technicon H-1 hematology instrument. Antagonism of the LTB₄-induced neutropenia response for the compounds tested is calculated.

Analgesic activity can be demonstrated e.g. in the Randall-Selitto test for analgesia, e.g. as described in Arch. Int. Pharmacodyn. Ther. 111, 409 (1957).

Antiinflammatory activity can be demonstrated by measuring the inhibition of the edema and inhibition of the influx of polymorphonuclear (PMN's) and mononuclear leukocytes (monocytes and macrophages) after oral administration in the rat model in which pleurisy is first induced by injecting carrageenin into the pleural cavity, e.g. according to A.P. Almeida et al., J. Pharmacol. Exp. Therap. 214, 74 (1980).

Bronchial effects such as anti-asthmatic activity, can be demonstrated in the antigen-induced guinea pig

bronchoconstriction test, e.g. as described by Anderson et al, Br. J. Pharmacol. 1983, 78, 67-74.

The trinitrobenzenesulfonic acid-induced chronic colitis test in the rat, e.g. as described by Wallace et al, Gastroenterology 1989, 96, 29-36, can be used to evaluate compounds for effects indicative of utility in inflammatory bowel diseases.

5 The arachidonic acid-induced mouse ear edema test, e.g. as described by Young et al, J. Invest. Dermatol. 1984, 82, 367-371 can be used to evaluate compounds for effects indicative of utility in dermatological disorders such as psoriasis.

The invention especially relates to compounds of formula I and pharmaceutically acceptable salts thereof, in which:


10 R is hydrogen, (C₁-C₁₂)alkoxycarbonyl or (C₂-C₁₂)alkenylloxycarbonyl, each of which is unsubstituted or substituted by phenyl, naphthyl, indanyl or fluorenyl, or is (C₃-C₇)cycloalkoxycarbonyl being unsubstituted or mono- or polysubstituted by lower alkyl, or is lower alkanoyl or phenyl-lower alkanoyl, or is benzoyl, naphthoyl, indanoyl or fluorenyl, or is (C₁-C₇)alkanesulfonyl, phenyl-(C₁-C₇)alkanesulfonyl, (C₃-C₇)cycloalkanesulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by lower alkyl, phenyl-lower alkyl or phenyl;
 15 R₁ is lower alkyl, lower alkenyl, lower alkynyl, phenyl-lower alkyl, phenyl-lower alkenyl, phenyl-lower alkynyl, (C₃-C₇)cycloalkyl, (C₃-C₇)cycloalkenyl, (C₃-C₇)-cycloalkyl-lower alkyl or (C₃-C₇)-cycloalkenyl-lower alkyl;
 X₁ and X₃, independently of one another, are O or S;
 X₂ is lower alkylene, lower alkylene-phenylene-lower alkylene or lower alkylene-naphthylene-lower alkylene; wherein the phenyl rings of formula I may be, independently of one another, substituted by one or more substituents selected from halogen, trifluoromethyl, lower alkyl, lower alkenyl, lower alkynyl, hydroxy, lower alkoxy,
 20 lower alkenyloxy, phenyl-lower alkoxy, cyclohexyl-lower alkoxy, lower alkanoyloxy, lower alkenoyloxy and phenyl-lower alkanoyloxy;
 wherein the aromatic radicals in the above definitions may be, independently of one another, substituted by one or more substituents selected from halogen, trifluoromethyl, lower alkyl, lower alkenyl, lower alkynyl, hydroxy, lower alkoxy, lower alkenyloxy, phenyl-lower alkoxy, cyclohexyl-lower alkoxy, lower alkanoyloxy, lower
 25 alkenoyloxy and phenyl-lower alkanoyloxy.

The invention especially relates to compounds of formula I and pharmaceutically acceptable salts thereof, in which:

30 R is hydrogen, C₁-C₁₂alkoxy-carbonyl, C₂-C₆alkanoyl, phenyl-C₂-C₆alkanoyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C₁-C₇alkyl, or C₁-C₇alkoxy, C₃-C₆cycloalkylcarbonyl which is unsubstituted or substituted by C₁-C₇alkyl, or is benzoyl, naphthoyl, indanoyl or fluorenyl, or is C₁-C₇alkanesulfonyl, phenyl-C₁-C₇alkanesulfonyl, C₃-C₇cycloalkanesulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by C₁-C₇alkyl, phenyl-C₁-C₇alkyl or phenyl;
 R₁ is C₁-C₇alkyl, phenyl-C₁-C₇alkyl, C₃-C₆cycloalkyl, and C₃-C₆cycloalkyl-lower alkyl;
 35 X₁ and X₃ each are -O-, or furthermore are, independently of one another, -O- or -S-;
 X₂ is C₂-C₇alkylene or C₂-C₄alkylene-phenylene-C₂-C₄alkylene;
 wherein phenyl in the above definitions is unsubstituted or, furthermore, independently of one another, substituted by a substituent selected from halogen, trifluoromethyl, C₁-C₇alkyl, and C₁-C₇alkoxy.

40 The invention especially relates to compounds of formula I and pharmaceutically acceptable salts thereof, in which -O-R₁ is located in position 2 (ortho) of the corresponding phenyl ring with respect to -X₁-; and -C(=NH)-NH-R is located in position 4 (para) of the corresponding phenyl ring with respect to -X₃-.

The invention especially relates to compounds of formula IA



and pharmaceutically acceptable salts thereof, in which:

55 R is hydrogen, C₁-C₁₂alkoxycarbonyl, such as methoxycarbonyl or octyloxycarbonyl, phenyl-C₁-C₄alkoxycarbonyl, such as benzyloxycarbonyl, C₂-C₆alkanoyl, such as acetyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C₁-C₄alkyl, or C₁-C₄alkoxy, such as 3,4-dimethoxybenzoyl, C₃-C₆cycloalkylcarbonyl which is unsubstituted or substituted by C₁-C₄alkyl, such as 2-isopropyl-5-methyl-cyclohexylcarbonyl;

R₁ is C₁-C₄-alkyl, such as methyl, ethyl or isopropyl, phenyl-C₁-C₄-alkyl, such as benzyl, C₃-C₆-cycloalkyl, such as cyclohexyl;

R₂, R₃, R₄ and R₆, independently of one another, are hydrogen, halogen, such as fluoro, chloro or bromo, C₁-C₄-alkoxy, such as methoxy, or C₁-C₄-alkyl, such as methyl, phenyl-C₁-C₄-alkoxy, such as benzyloxy, or C₃-C₆-cycloalkyl-C₁-C₄-alkoxy, such as cyclohexylmethoxy;

X₁ and X₃ are -O-;

X₂ is C₄-C₇-alkylene, such as pentylenes, or C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene, such as 1,2-ethylene-1,4-phenylene-1,2-ethylene.

The invention especially relates to compounds of formula IA and pharmaceutically acceptable salts thereof, in which:

R is hydrogen;

R₁ is C₁-C₄-alkyl, such as methyl or isopropyl, phenyl-C₁-C₄-alkyl, such as benzyl, C₃-C₆-cycloalkyl, such as cyclohexyl;

R₂, R₃, R₄ and R₆, independently of one another, are hydrogen, halogen, such as fluoro, chloro or bromo, C₁-C₄-alkoxy, such as methoxy, or C₁-C₄-alkyl, such as methyl;

X₁ and X₃ are -O-;

X₂ is C₄-C₇-alkylene, such as pentylenes, or C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene, such as 1,2-ethylene-1,4-phenylene-1,2-ethylene.

The invention especially relates to compounds of formula IA and pharmaceutically acceptable salts thereof, in which:

R is C₁-C₄-alkanesulfonyl, such as methane-, ethane- or isopropanesulfonyl, phenyl-C₁-C₄-alkanesulfonyl, such as benzyulsulfonyl, C₃-C₇-cycloalkane-sulfonyl, such as cyclohexanesulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by C₁-C₄-alkyl, phenyl-C₁-C₄-alkyl or phenyl;

R₁ is C₁-C₄-alkyl, such as methyl or isopropyl, phenyl-C₁-C₄-alkyl, such as benzyl, C₃-C₆-cycloalkyl, such as cyclohexyl;

R₂, R₃, R₄ and R₆, independently of one another, are hydrogen, halogen, such as fluoro, chloro or bromo, C₁-C₄-alkoxy, such as methoxy, or C₁-C₄-alkyl, such as methyl;

X₁ and X₃ are -O-;

X₂ is C₄-C₇-alkylene, such as pentylenes, or C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene, such as 1,2-ethylene-1,4-phenylene-1,2-ethylene.

The invention further especially relates to compounds of formula IA and pharmaceutically acceptable salts thereof, in which:

R is C₁-C₁₂-alkoxycarbonyl, such as methoxycarbonyl or octyloxycarbonyl, phenyl-C₁-C₄-alkoxycarbonyl, such as benzyloxycarbonyl, C₂-C₆-alkanoyl, such as acetyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C₁-C₄-alkyl, or C₁-C₄-alkoxy, such as 3,4-dimethoxybenzoyl, C₃-C₆-cycloalkylcarbonyl which is unsubstituted or substituted by C₁-C₄-alkyl, such as 2-isopropyl-5-methyl-cyclohexylcarbonyl;

R₁ is C₁-C₄-alkyl, such as methyl or isopropyl, or phenyl-C₁-C₄-alkyl, such as benzyl;

R₂ is hydrogen or halogen, such as fluoro, C₁-C₄-alkoxy, such as methoxy;

R₃ is hydrogen or halogen, such fluoro;

R₄ and R₆, independently of one another, are hydrogen, halogen, such as fluoro, or C₁-C₄-alkoxy, such as methoxy;

X₁ and X₃ are -O-;

X₂ is C₄-C₇-alkylene, especially pentylenes.

The invention further especially relates to compounds of formula IA and pharmaceutically acceptable salts thereof, in which:

R is hydrogen;

R₁ is C₁-C₄-alkyl, such as methyl or isopropyl, or phenyl-C₁-C₄-alkyl, such as benzyl;

R₂ is hydrogen or halogen, such as fluoro, C₁-C₄-alkoxy, such as methoxy;


R₃ is hydrogen or halogen, such fluoro;

R₄ and R₆, independently of one another, are hydrogen, halogen, such as fluoro, or C₁-C₄-alkoxy, such as methoxy;

X₁ and X₃ are -O-;

X₂ is C₄-C₇-alkylene, especially pentylenes.

The invention further especially relates to compounds of formula IB



10

and pharmaceutically acceptable salts thereof, in which:

R is hydrogen;

R₁ is C₁-C₄-alkyl, such as methyl or isopropyl;

R₂ is hydrogen, halogen, such as fluoro, or C₁-C₄-alkoxy, such as methoxy;

15 R₃ is hydrogen or halogen, such as fluoro;

X₁ and X₃ are -O-;

X₂ is C₄-C₇-alkylene, especially pentylene.

The invention further especially relates to compounds of formula (IB) and pharmaceutically acceptable salts thereof, in which:

20

R is hydrogen;

R₁ is C₁-C₄-alkyl, such as methyl or isopropyl;

R₂ and R₃, independently of one another, are hydrogen or halogen, such as fluoro;

X₁ and X₃ are -O-;

X₂ is C₄-C₇-alkylene, especially pentylene.

25

Preferred embodiments of the invention relate to the above compounds of formula IA and IB wherein at least one of R₂, R₃, R₄ and R₅ is hydrogen. Preferred are the above compounds wherein R₅ is hydrogen, and further wherein R₄ and R₆ are hydrogen.

The invention relates in particular to the novel compounds itemized in the examples and to the manners of preparation described therein.

30


The invention further relates to methods for the preparation of the compounds according to the invention.

The preparation of compounds of the formula I is, for example, characterized in that,

a) for the manufacture of compounds of the formula I in which R is hydrogen, in a compound of the formula IIa

35

40




45

or a salt thereof in which Z₁ is a radical which can be converted into the variable -C(=NH)-NH-R, Z₁ is converted into the variable -C(=NH)-NH-R, or


b) a compound of the formula IIIa

50



55

or a salt thereof is reacted with a compound of the formula IIIb



10 or a salt thereof in which Z_2 is a group of the formula $-X_1-X_2-Z_4$ and Z_3 is $-Z_5$, or Z_2 is $-Z_5$ and Z_3 is a group of the formula $Z_5-X_2-X_3$, wherein one of the radicals Z_4 and Z_5 is hydroxy or mercapto and the other is hydroxy, mercapto or reactive esterified hydroxy,

and, if desired, a compound of the formula I or a salt thereof obtainable according to the process or in another manner is converted into another compound or a salt thereof according to the invention, a free compound of the formula I obtainable according to the process is converted into a salt, a salt obtainable according to the process is converted into the free compound of the formula I or into another salt, or a mixture of isomers obtainable according to the process is resolved and the desired compound is isolated.

Salts of starting materials which contain at least one basic centre, for example of the formula IIa, are appropriate acid addition salts, while salts of starting materials which contain an acid group are present as salts with bases.

20 A radical Z_1 which can be converted into the variable $-C(=NH)-NH-R$ is, for example, (lower) alkoxy-iminocarbonyl or halogeno-iminocarbonyl [Halogeno- $C(=NH)-$].

Reactive esterified hydroxy (e.g. Z_4 or Z_5) is, in particular, hydroxy esterified with a strong inorganic acid or organic sulfonic acid, and is, for example, halogen, such as chlorine, bromine or iodine, sulfonyloxy, such as hydroxysulfonyloxy, halosulfonyloxy, such as, fluorosulfonyloxy, (C_1-C_7) alkanesulfonyloxy which, if desired, is substituted, for example, by halogen, such as, methane- or trifluoromethanesulfonyloxy, (C_6-C_7) cycloalkanesulfonyloxy, such as, cyclohexanesulfonyloxy, or benzenesulfonyloxy which, if desired, is substituted, for example by (C_1-C_7) alkyl or halogen, such as, p-bromobenzene- or p-toluenesulfonyloxy.

The reactions described in the variants above and below are carried out in a manner known per se, for example in the absence or in the customary manner in the presence of a suitable solvent or diluent or a mixture thereof, the reaction being carried out, according to need, with cooling, at room temperature or with warming, for example in a temperature range from about -80°C up to the boiling point of the reaction medium, preferably from about -10°C to about $+180^\circ\text{C}$, and, if necessary, in a closed vessel, under pressure, in an inert gas atmosphere and/or under anhydrous conditions.

35 Process a):

Alkoxy-iminocarbonyl is, for example C_1-C_4 alkoxy-iminocarbonyl such as methoxy- or ethoxy-iminocarbonyl, whereas halogeno-iminocarbonyl is, for example chloro-iminocarbonyl.

40 Preferably, those compounds of the formula IIa are employed in which Z_1 is C_1-C_4 alkoxy-iminocarbonyl such as methoxy- or ethoxy-iminocarbonyl. The reaction is preferably carried out by reacting with ammonia and using an acid resulting in the corresponding acid addition salt of the product. As acids are used, for example, inorganic acids, such as mineral acids, for example sulfuric acid, a phosphoric or hydrohalic acid, or with organic carboxylic acids, such as (C_1-C_4) alkanecarboxylic acids which, for example, are unsubstituted or substituted by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, 45 malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, for example aspartic or glutamic acid, benzoic acid or with organic sulfonic acids, such as (C_1-C_4) alkane- or arylsulfonic acids which are unsubstituted or substituted, for example, by halogen, for example methane- or toluenesulfonic acid. Preferred acids are hydrohalic acids, especially hydrochloric acid, organic sulfonic acids, especially methanesulfonic acid, or dicarboxylic acids, especially maleic acid.

Process b):

55 Preferably, those compounds of the formulae IIIa and IIIb are employed in which Z_2 is a group of the formula $-X_1-X_2-Z_4$, wherein Z_4 is reactive esterified hydroxy, preferably halogen, especially bromine, and Z_3 is hydroxy or mercapto.


The reaction is carried out preferably in the presence of a base.

Suitable bases are, for example, alkali metal hydroxides, hydrides, amides, alkanolates, carbonates, tri-


phenylmethyldes, di(lower alkyl)amides, aminoalkylamides or lower alkyl silylamides, or naphthaleneamines, lower alkylamines, basic heterocycles, ammonium hydroxides, and also carbocyclic amines. Examples which may be mentioned are sodium hydroxide, sodium hydride, sodium amide, sodium (m)ethoxide, potassium tert-butoxide, potassium carbonate, lithium triphenylmethyldes, lithium diisopropylamide, potassium 3-(aminopropyl)amide, potassiumbis(trimethylsilyl)amide, dimethylaminonaphthalene, di- or triethylamine, or ethyldiisopropylamine, N-methylpiperidine, pyridine, benzyltrimethylammonium hydroxide, 1,5-diazabicyclo[4.3.0]non-5-ene (DBN) and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU). Preferred is cesium carbonate.

The starting material can be prepared following methods known per se.

In order to prepare the starting material of the formula IIa, for example, a compound of the formula




in which Z_2 preferably is a group of the formula $-X_1-X_2-Z_4$, wherein Z_4 preferably is reactive esterified hydroxy, is reacted with a compound of the formula



in which Z_3 preferably is hydroxy or mercapto, following the method as described in process b).

If one of variables Z_2 and Z_4 represents reactive esterified hydroxy, the other preferably represents hydroxy or mercapto.

A compound of the formula IIIb can be obtained, for example, by converting Z_1 of a compound of the formula (IIc) in which Z_1 is a radical which can be converted into the variable $-C(=NH)-NH-R$ following the method as described in process a). Compounds of formulae (IIb) and (IIc) are known or can be prepared according to methods known per se. Preferably, Z_2 is a group of the formula $-X_1-X_2-Z_4$, wherein Z_4 is preferably reactive esterified hydroxy, such as chlorine or bromine, and Z_3 is hydroxy or mercapto. A corresponding compound of the formula IIc, wherein Z_2 is a group of the formula $-X_1-X_2-Z_4$ and Z_4 is preferably reactive esterified hydroxy, such as chlorine or bromine, can be obtained, for example, by reacting a compound of the formula $Z_4-X_2-Z_4$ (IIId) with a compound of the formula



preferably in the presence of a base.

The invention further includes any variant of the present processes, in which an intermediate product obtainable at any stage thereof is used as starting material and the remaining steps are carried out, or the process is discontinued at any stage thereof, or in which the starting materials are formed under the reaction conditions, or in which the reaction components are used in the form of their salts or optically pure antipodes.

A compound according to the invention which is obtainable by the process can be converted into another compound according to the invention in a manner known per se.

If R is hydrogen, the corresponding amidino group can be N-acylated accordingly. The acylation is carried

out in a manner known per se using a suitable acylating agent. An example of a suitable acylating agent is a compound of the formula Ac-Z_6 (IIia), where Ac denotes an acyl radical corresponding to the variable R, and Z_6 denotes in particular reactive activated hydroxyl. Appropriate hydroxyl can be activated, for example, by strong acids such as hydrohalic or carboxylic acid, for example by hydrochloric, hydrobromic acid, an optionally substituted, for example by halogen, alkanecarboxylic acid or by an acid of the formula Ac-OH , or by suitable activating or coupling reagents of the type detailed hereinafter, in particular in situ. Ac-Z_6 can furthermore represent an activated ester, where Z_6 denotes, in particular, cyanomethoxy, (4-)nitrophenoxy or polyhalogeno-, such as pentachloro-, -phenoxy. Activating and coupling reagents which can be employed are, in particular, carbodiimides, for example N,N'-di-C₁-C₄-alkyl- or N,N'-di-C₆-C₇-cycloalkyl-carbodiimide, such as diisopropylcarbodiimide or N,N'-dicyclohexylcarbodiimide, advantageously with the addition of an activating catalyst such as N-hydroxysuccinimide or optionally substituted, for example by halogen, C₁-C₇-alkyl or C₁-C₇-alkoxy, N-hydroxy-benzotriazole or N-hydroxy-5-norbornene-2,3-dicarboxamide, furthermore C₁-C₄-alkyl halogenoformate, for example isobutyl chloroformate, suitable carbonyl compounds, for example N,N-carbonyldiimidazole, suitable 1,2-oxazolium compounds, for example 2-ethyl-5-phenyl-1,2-oxazolium 3'-sulfonate or 2-tert-butyl-5-methyl-isoxazolium perchlorate, suitable acylamino compounds, for example 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline, or suitable phosphoryl cyanamides or azides, for example diethylphosphoryl cyanamide or diphenylphosphoryl azide, furthermore triphenylphosphine disulfide or 1-C₁-C₄-alkyl-2-halogeno-pyridinium halides, for example 1-methyl-2-chloropyridinium iodide. Z_6 preferably denotes halogen such as chlorine or bromine, and Ac-O .

If the compounds of the formula (I) or (IA) or (IB) contain unsaturated radicals, such as (lower)alkenyl groups, these can be converted into saturated radicals in a manner known per se. Thus, for example, multiple bonds are hydrogenated by catalytic hydrogenation in the presence of hydrogenation catalysts, suitable for this purpose being, for example, nickel, such as Raney nickel, and noble metals or their derivatives, for example oxides, such as palladium or platinum oxide, which may be applied, if desired, to support materials, for example to carbon or calcium carbonate. The hydrogenation may preferably be carried out at pressures between 1 and about 100 at and at room temperature between about -80° to about 200°C, in particular between room temperature and about 100°C. The reaction is advantageously carried out in a solvent, such as water, a lower alkanol, for example ethanol, isopropanol or n-butanol, an ether, for example dioxane, or a lower alkanecarboxylic acid, for example acetic acid.

The invention also relates to any novel starting materials and processes for their manufacture and their use.

Depending on the choice of starting materials and methods, the new compounds may be in the form of one of the possible isomers or mixtures thereof, for example, as substantially pure geometric (cis or trans) isomers, optical isomers (antipodes), racemates, or mixtures thereof. The aforesaid possible isomers or mixtures thereof are within the purview of this invention.

Any resulting mixtures of isomers can be separated on the basis of the physico-chemical differences of the constituents, into the pure geometric or optical isomers, diastereoisomers, racemates, for example by chromatography and/or fractional crystallization.

Any resulting racemates of final products or intermediates can be resolved into the optical antipodes by known methods, e.g. by separation of the diastereoisomeric salts thereof, obtained with an optically active acid or base, and liberating the optically active acidic or basic compound. The amidine (wherein R_3 represents hydrogen) can thus be resolved into their optical antipodes e.g. by fractional crystallization of a salt formed with an optically active acid.

Finally, the compounds of the invention are either obtained in the free form, or as a salt thereof.

In view of the close relationship between the free compounds and the compounds in the form of their salts, whenever a compound is referred to in this context, a corresponding salt is also intended, provided such is possible or appropriate under the circumstances.

The compounds, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for their crystallization.

The pharmaceutical compositions according to the invention are those suitable for enteral, such as oral or rectal, transdermal and parenteral administration to mammals, including man, to antagonize LTB_4 receptors, and for the treatment of a condition or syndrome responsive to the selective antagonism of LTB_4 receptors, comprising an effective amount of a pharmacologically active compound of the invention, alone or in combination, with one or more pharmaceutically acceptable carriers.

The novel pharmaceutical products contain, for example, from about 10 % to about 80 %, preferably from about 20 % to about 60 %, of the active compound. Examples of pharmaceutical products according to the invention for enteral or parenteral administration are those in dose-unit forms such as coated tablets, tablets, capsules or suppositories, as well as ampoules. These are prepared in a manner known per se, for example

using conventional mixing, granulating, coating, dissolving or freeze-drying processes. Thus, pharmaceutical products for oral use can be obtained by combining the active compound with solid excipients, where appropriate granulating a mixture which is obtained, and processing the mixture or granules, if desired or necessary, after addition of suitable auxiliaries to tablets or cores of coated tablets.

5 The pharmacologically active compounds of the invention are useful in the manufacture of pharmaceutical compositions comprising an effective amount thereof in conjunction or admixture with excipients or carriers suitable for either enteral or parenteral application. Preferred are tablets and gelatin capsules comprising the active ingredient together with a) diluents, e.g. lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine; b) lubricants, e.g. silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethyleneglycol; for tablets
10 also c) binders e.g. magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and/or polyvinylpyrrolidone; if desired d) disintegrants, e.g. starches, agar, alginic acid or its sodium salt, or effervescent mixtures; and/or e) absorbants, colorants, flavors and sweeteners. Injectable compositions are preferably aqueous isotonic solutions or suspensions, and suppositories are advantageously prepared from fatty emulsions or suspensions. Said compositions may be sterilized and/or contain adjuvants,
15 such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. Cores of coated tablets are provided with suitable, optionally enteric, coatings, using, inter alia, concentrated sugar solutions which optionally contain gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions in suitable organic solvents or solvent mixtures or, for the preparation of enteric coatings, solutions of suitable cellulose products such as acetyl cellulose phthalate or hydroxypropylmethylcellulose phthalate. Colorants or pigments can be added to the tablets or coatings
20 of coated tablets, for example, to identify or to indicate various doses of active compound. In addition, they may also contain other therapeutically valuable substances. Said compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1 to 75 %, preferably about 1 to 50 %, of the active ingredient.

25 Suitable formulations for transdermal application include an effective amount of a compound of the invention with carrier. Advantageous carriers include absorbable pharmacologically acceptable solvents to assist passage through the skin of the host. Characteristically, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the compound of the skin of the host at a controlled and predetermined rate over a
30 prolonged period of time, and means to secure the device to the skin.

In conjunction with another active ingredient, a compound of the invention may be administered either simultaneously, before or after the other active ingredient, either separately by the same or different route of administration or together in the same pharmaceutical formulation.


35 The invention further particularly relates to a method of the treatment of a condition or syndrome responsive to the selective antagonism of LTB_4 receptors, such as rheumatoid arthritis, inflammatory bowel disease, psoriasis, non-steroidal-antiinflammatory-drug-induced gastropathy, adult respiratory distress syndrome (ARDS), myocardial infarction, allergic rhinitis, hemodialysis-induced neutropenia, and late phase asthma.

The dosage of active compound administered is dependent on the species of warm-blooded animal (mammal), the body weight, age and individual condition, and on the form of administration. A unit dosage for oral
40 administration to a mammal of about 70 kg may contain e.g. between about 1 and about 1000 mg/kg per day of the active ingredient.

The following examples are intended to illustrate the invention and are not to be construed as being limitations thereon. Temperatures are given in degrees Centigrade. If not mentioned otherwise, all evaporations are performed under reduced pressure, preferably between about 15 and 100 mm Hg. The structure of final
45 products, intermediates and starting materials is confirmed by standard analytical methods, e.g. microanalysis and spectroscopic characteristics (e.g. MS, IR, NMR). Abbreviations used are those conventional in the art.

Example 1:

50 To a suspension of 9.3 g (23.6 mmol) of ethyl 4-[5-(2-methoxyphenoxy)pentoxy] benzenecarboximidate hydrochloride in 230 mL of ethanol cooled in an ice-bath is bubbled ammonia gas for 15 minutes. The resulting solution is stirred at room temperature overnight. The solvent is removed under reduced pressure and the residue dissolved in methanol and an equivalent of methanesulfonic acid added and the solution diluted with a large volume of ether. The crude salt produced is recrystallized from ethanol to give 4-[5-(2-methoxyphenoxy)pentoxy] benzenecarboximidamide monomethanesulfonate, m.p. 115-117°C.
55



Similarly prepared are the hydrochloride salt, m.p. 89-90°C, and the maleate salt, m.p. 163-165°C.

The starting material can be prepared, for example as follows.

To a solution of 16.9 g (135 mmol) p-cyanophenol in 169 mL of dimethylformamide (DMF), cooled in an ice-bath is added in portions 6.5 g (135 mmol) sodium hydride (50% dispersion in oil). The reaction is stirred and allowed to warm-up to room temperature until no further hydrogen evolution is observed. The solution is then cooled in an ice-bath and a solution of 1-bromo-5-chloro-pentane 25 g (135 mmol) in 84 mL of dimethylformamide is added. The reaction is allowed to warm up to room temperature and stirred for 18 h. The reaction is then poured onto water and the resulting solid filtered off and dried to give 4-(5-chloropentoxy) benzonitrile, m.p. 52-54°C.

To a solution of 32.7 g (146 mmol) of 4-(5-chloropentoxy) benzonitrile in 350 mL acetone is added 38.3 g (150 mmol) of sodium iodide in 230 mL of acetone. The reaction is heated to reflux and stirred overnight. The acetone is removed under reduced pressure and the residue partitioned between water and ether. The ethereal layer is dried, filtered and evaporated to give 4-(5-iodopentoxy) benzonitrile, m.p. 51-54°C.

To a solution of 5.1 g (41 mmol) of gualacol in 41 mL of DMF, cooled in an icebath, is added in portions, 20 g (41 mmole) of sodium hydride (50% dispersion in oil). The reaction is allowed to warm up to room temperature and stirred until no further hydrogen evolution is observed. The solution is then again cooled in an ice-bath and a solution of 13 g (41 mmol) 4-(5-iodopentoxy) benzonitrile in 82 mL DMF added. The reaction is allowed to warm up to room temperature and stirred for 18 h before pouring the mixture into water. The resulting solid is filtered off and dried to give 4-[5-(2-methoxyphenoxy)pentoxy] benzonitrile, m.p. 61-63°C.

To a solution of 11.8 g (38 mmol) 4-[5-(2-methoxyphenoxy)pentoxy] benzonitrile in a mixture of 161 mL of methylene chloride and 6.8 mL of ethanol is bubbled hydrogen chloride gas for a period of 15 min. The reaction is then stirred at room temperature for 6 days. The solvents are removed under reduced pressure and the residue triturated with ether to give ethyl 4-[5-(2-methoxyphenoxy)pentoxy] benzenecarboximidate hydrochloride, m.p. 114-116°C.

Example 2:

In a way analogously as described in Example 1, the following compounds can be manufactured:

- 4-[5-(2,3-dimethoxyphenoxy)pentoxy]benzenecarboximidamidesulfonate, m.p. = 163-165°C.
- 4-[5-(2,6-dimethoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 123-126°C.
- 4-[5-(2-(1-methylethoxy)phenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 102-107°C.
- 4-[5-(2-(phenylmethoxy)phenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 116-123°C.
- 4-[5-(2-ethoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 128-129°C.
- 4-[5-(2-hexyloxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 129-130°C.
- 4-[5-(5-fluoro-2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 130-131°C.
- 4-[5-(4,5-difluoro-2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. 140-142°C.
- 4-[5-(4,5-difluoro-2-(1-methylethoxy)phenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 136-137°C.
- 3-methoxy-4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 140°C.
- 3-fluoro-4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 130°C.
- 3,5-dichloro-4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 124-126°C.
- 3-chloro-4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 148-149.5°C.
- 2-fluoro-4-[5-(2-(1-methylethoxy)phenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 142-144°C.
- 4-[4-(2-methoxyphenoxy)butoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 139-140°C.
- 4-[6-(2-methoxyphenoxy)hexyloxy]benzenecarboximidamide monomethanesulfonate, m.p. = 147-148°C.

4-[4-[(2-methoxyphenoxy)methyl]phenylmethoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 182-185°C.

4-[5-[5-fluoro-2-(1-methylethoxy)phenoxy]pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 94-96°C.

5 4-[5-(4-fluoro-2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 129-131°C.

4-[5-(4-fluoro-2-(1-methylethoxy)phenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. 124-127°C.

4-[5-[4-(1,1,3,3-tetramethylbutyl)-2-methoxyphenoxy]pentoxy]benzenecarboximidamide hydrochloride.

10

Example 3:

A stirred, 0°C solution of ethyl 4-[5-(2-phenylmethoxy)phenoxy]pentoxy benzenecarboximidoate (320 mg, 0.71 mmol) in 8 mL anhydrous ethanol in a pressure tube is treated with anhydrous ammonia for 15 minutes. The pressure tube is sealed and heated to 100°C for 45 minutes. Upon cooling and concentrating in vacuo, the resulting material is dissolved in methanol and an equivalent of methanesulfonic acid added and the solution diluted with a large volume of ether. The crude salt produced is recrystallized from ethanol to active 4-[5-[(2-phenylmethoxy)phenoxy]pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. 116-123°C.

The starting material, ethyl 4-[5-[(2-phenylmethoxy)phenoxy]pentoxy] benzenecarboximidoate, can be prepared, for example, as follows:

A stirred, 0°C solution of 4-hydroxybenzonitrile (50.7 g, 426 mmol) in 1400 mL of dichloromethane and 75 mL anhydrous ethanol is treated with anhydrous hydrogen chloride gas (110 g) over 1.5 hours. This solution is stirred at room temperature for 64 hours and the resulting solids collected and washed with 500 mL diethyl ether and 2 x 1000 mL of ethyl acetate. The remaining solids (60.4 g) were dissolved in 1200 mL of water and the residual solids filtered. To the filtrate is added a solution of sodium hydroxide (12.57 g) in 150 mL water. The resulting white solid is filtered and washed with water to afford ethyl 4-hydroxybenzenecarboximidoate.

A stirred solution of ethyl 4-hydroxybenzenecarboximidoate (32.0 g, 194 mmol) in 250 mL anhydrous N,N-dimethylformamide is treated with cesium carbonate (75.7 g, 232 mmol) and 1,5-dibromopentane (89.1 g, 387 mmol) and heated at 70° for 1.5 hours. After cooling to room temperature, the reaction is filtered and the resulting filtrate concentrated in vacuo to afford a yellow oil (85.7 g). This material is purified by chromatography on silica gel (850 g) with 10-60% ethyl acetate/hexanes as the eluent to afford ethyl 4-[5-bromopentoxy] benzenecarboximidoate as a colorless oil.

A stirred solution of 2-phenylmethoxyphenol (200 mg, 1 mmol) in 5 mL anhydrous N,N-dimethylformamide is treated with cesium carbonate (330 mg, 1 mmol) and ethyl 4-[5-bromopentoxy]benzenecarboximidoate (300 mg, 1 mmol) and stirred at R.T. for 24 h. The reaction is poured into water and the solids filtered off and dried in vacuo to afford ethyl 4-[5-[(2-phenylmethoxy)phenoxy]pentoxy]benzenecarboximidoate m.p. 77-79°C.

Example 4:

40 In a way analogous as described in Example 3, the following compounds can be manufactured:

4-[5-[(2-methoxy-4-(1-methylethoxy)phenoxy)]pentoxy]benzenecarboximidamide monohydrochloride.

4-[5-[(2-methoxy-4-(phenylmethoxy)phenoxy)]pentoxy]benzenecarboximidamide monohydrochloride.

4-[5-[4-(cyclohexylmethoxy)-2-methoxyphenoxy]pentoxy]benzenecarboximidamide monohydrochloride.

4-[5-[2-methoxy-4-(2-methylpropoxy)phenoxy]pentoxy]benzenecarboximidamide monohydrochloride.

45 4-[5-(2-methoxy-5-methylphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. 136-138°C.

4-[5-(4-bromo-2-methoxyphenoxy)pentoxy]benzenecarboximidamide monohydrochloride, m.p. = 85-86°C.

4-[5-[(2-methoxyphenyl)thio]pentoxy]benzenecarboximidamide monomethanesulfonate, m.p. = 125-127°C.

4-[5-(2-methoxyphenoxy)pentoxy]thio]benzenecarboximidamide monomethanesulfonate, m.p. = 95-98°C.

50 Example 5:

a) Preparation of 10,000 tablets each containing 20 mg of the active ingredient, for example, 4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate:

active ingredient	200.00 g
55 Lactose	2,535.00 g
Corn starch	125.00 g
Polyethylene glycol 6,000	150.00 g
Magnesium stearate	40.00 g

Purified water q.s.

Procedure:

5 All the powders are passed through a screen with openings of 0.6 mm. The drug substance, lactose, magnesium stearate and half of the starch are mixed in a suitable mixer. The other half of the starch is suspended in 65 ml of water and the suspension added to the boiling solution of the polyethylene glycol in 250 ml of water. The paste formed is added to the powders, which are granulated, if necessary, with an additional amount of water. The granulate is dried overnight at 35°C broken on a screen with 1.2 mm openings and compressed into tablets, using concave punches uppers bisected.

10 Analogously tablets are prepared, containing about 10-100 mg of one of the other compounds disclosed and exemplified herein.

b) Preparation of 1,000 capsules each containing 40 mg of the active ingredient, for example, 4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide monomethanesulfonate:

15 active ingredient	40.00 g
Lactose	177.00 g
Modified starch	80.00 g
Magnesium stearate	3.00 g


20 Procedure:

All the powders are passed through a screen with openings of 0.6 mm. The drug substance is placed in a suitable mixer and mixed first with the magnesium stearate, then with the lactose and starch until homogenous. No. 2 hard gelatin capsules are filled with 300 mg of said mixture each, using a capsule filling machine.

25 Analogously capsules are prepared, containing about 10-100 mg of the other compounds disclosed and exemplified herein.

Claims

30 1. A compound of the formula I



wherein:

R is hydrogen or an acyl radical which is derived from an organic carbonic acid, an organic carboxylic acid, a sulfonic acid, or a carbamic acid;

R₁ is a substituent selected from an aliphatic hydrocarbon radical, an araliphatic hydrocarbon radical and a cycloaliphatic hydrocarbon radical;

X₁ and X₃, independently of one another, are oxygen (-O-) or sulphur (-S-); and

X₂ is a divalent aliphatic hydrocarbon radical which may be interrupted by an aromatic radical;


50 wherein the phenyl rings of formula I may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid;

55 wherein aryl in the above definitions may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid; and

wherein a cycloaliphatic hydrocarbon radical may be substituted by an aliphatic radical; or a pharmaceut-

ically acceptable salt thereof.

2. A compound according to claim 1 wherein R is hydrogen, (C₁-C₁₂)-alkoxycarbonyl or (C₂-C₁₂)-alkenyloxycarbonyl, each of which is unsubstituted or substituted by phenyl, naphthyl, indanyl or fluorenyl, or is (C₃-C₇)-cycloalkoxycarbonyl being unsubstituted or mono- or polysubstituted by lower alkyl, or is lower alkanoyl or phenyl-lower alkanoyl, or is benzoyl, naphthoyl, indanoyl or fluorenyl, or is (C₁-C₇)-alkanesulfonyl, phenyl-(C₁-C₇)-alkanesulfonyl, (C₃-C₇)-cycloalkanesulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by lower alkyl, phenyl-lower alkyl or phenyl;
 5 R₁ is lower alkyl, lower alkenyl, lower alkynyl, phenyl-lower alkyl, phenyl-lower alkenyl, phenyl-lower alkynyl, (C₃-C₇)-cycloalkyl, (C₃-C₇)-cycloalkenyl, (C₃-C₇)-cycloalkyl-lower alkyl or (C₃-C₇)-cycloalkenyl-lower alkyl;
 X₁ and X₃, independently of one another, are O or S;
 X₂ is lower alkylene, lower alkylene-phenylene-lower alkylene or lower alkylene-naphthylene-lower alkylene;
 15 wherein the phenyl rings of formula I may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, lower alkyl, lower alkenyl, lower alkynyl, hydroxy, lower alkoxy, lower alkenyloxy, phenyl-lower alkoxy, lower alkanoyloxy, lower alkenoyloxy and phenyl-lower alkanoyloxy;
 wherein the aromatic radicals in the above definitions may be, independently of one another, substituted
 20 by one or more substituents selected from halogen, trifluoromethyl, lower alkyl, lower alkenyl, lower alkynyl, hydroxy, lower alkoxy, lower alkenyloxy, phenyl-lower alkoxy, cyclohexyl-lower alkoxy, lower alkanoyloxy, lower alkenoyloxy and phenyl-lower alkanoyloxy, or a pharmaceutically acceptable salt thereof.
3. A compound according to claim 1 wherein R is hydrogen, C₁-C₁₂-alkoxy-carbonyl, C₂-C₆-alkanoyl, phenyl-C₂-C₆-alkanoyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C₁-C₇-alkyl, or C₁-C₇-alkoxy, C₃-C₆-cycloalkylcarbonyl which is unsubstituted or substituted by C₁-C₇-alkyl, or is benzoyl, naphthoyl, indanoyl or fluorenyl, or is C₁-C₇-alkanesulfonyl, phenyl-C₁-C₇-alkanesulfonyl, C₃-C₇-cycloalkanesulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by C₁-C₇-alkyl, phenyl-C₁-C₇-alkyl or phenyl;
 30 R₁ is C₁-C₇-alkyl, phenyl-C₁-C₇-alkyl or C₃-C₆-cycloalkyl;
 X₁ and X₃ each are -O-, or furthermore are, independently of one another, -O- or -S-;
 X₂ is C₂-C₇-alkylene or C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene;
 wherein phenyl in the above definitions is unsubstituted or, furthermore, independently of one another, substituted by one or more substituents selected from halogen,
 35 trifluoromethyl, C₁-C₇-alkyl, and C₁-C₇-alkoxy; or a pharmaceutically acceptable salt thereof.
4. A compound according to claim 1 wherein -O-R₁ is located in position 2 (ortho) of the corresponding phenyl ring with respect to -X₁-; and -C(=NH)-NH-R is located in position 4 (para) of the corresponding phenyl ring with respect to -X₃-; or a pharmaceutically acceptable salt thereof.
5. A compound according to claim 1 of formula IA



wherein R is hydrogen, C₁-C₁₂-alkoxycarbonyl, phenyl-C₁-C₄-alkoxycarbonyl, C₂-C₆-alkanoyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C₁-C₄-alkyl, or C₁-C₄-alkoxy, C₃-C₆-cycloalkylcarbonyl which is unsubstituted or substituted by C₁-C₄-alkyl;

R₁ is C₁-C₄-alkyl, phenyl-C₁-C₄-alkyl, C₃-C₆-cycloalkyl;


R₂, R₃, R₄ and R₅, independently of one another, are hydrogen, halogen, C₁-C₄-alkoxy, or C₁-C₄-alkyl;

X₁ and X₃ are -O-;

X_2 is C_4 - C_7 -alkylene or C_2 - C_4 -alkylene-phenylene- C_2 - C_4 -alkylene; or a pharmaceutically acceptable salt thereof.

6. A compound according to claim 5 of formula IA wherein R is hydrogen;
 R_1 is C_1 - C_4 -alkyl, phenyl- C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkyl;
 R_2 , R_3 , R_4 and R_6 , independently of one another, are hydrogen, halogen, C_1 - C_4 -alkoxy, or C_1 - C_4 -alkyl;
 X_1 and X_3 are -O-;
 X_2 is C_4 - C_7 -alkylene or C_2 - C_4 -alkylene-phenylene- C_2 - C_4 -alkylene; or a pharmaceutically acceptable salt thereof.
7. A compound according to claim 5 of formula IA wherein R is C_1 - C_4 -alkanesulfonyl, phenyl- C_1 - C_4 -alkanesulfonyl, C_3 - C_7 -cycloalkane-sulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by C_1 - C_4 -alkyl, phenyl- C_1 - C_4 -alkyl or phenyl;
 R_1 is C_1 - C_4 -alkyl, phenyl- C_1 - C_4 -alkyl, C_3 - C_6 -cycloalkyl;
 R_2 , R_3 , R_4 and R_6 , independently of one another, are hydrogen, halogen, C_1 - C_4 -alkoxy or C_1 - C_4 -alkyl;
 X_1 and X_3 are -O-;
 X_2 is C_4 - C_7 -alkylene or C_2 - C_4 -alkylene-phenylene- C_2 - C_4 -alkylene; or a pharmaceutically acceptable salt thereof.
8. A compound according to claim 5 of formula IA wherein
R is hydrogen;
 R_1 is C_1 - C_4 -alkyl or phenyl- C_1 - C_4 -alkyl;
 R_2 is hydrogen or C_1 - C_4 -alkoxy;
 R_3 is hydrogen or halogen;
 R_4 and R_6 , independently of one another are hydrogen, halogen or C_1 - C_4 -alkoxy;
 X_1 and X_3 are -O-;
 X_2 is C_4 - C_7 -alkylene; or a pharmaceutically acceptable salt thereof.
9. A compound according to claim 8 wherein R_4 and R_6 are hydrogen.
10. A compound according to claim 5 of formula IA wherein R is C_1 - C_{12} -alkoxycarbonyl, phenyl- C_1 - C_4 -alkoxycarbonyl, C_2 - C_6 -alkanoyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C_1 - C_4 -alkyl, or C_1 - C_4 -alkoxy, C_3 - C_6 -cycloalkylcarbonyl which is unsubstituted or substituted by C_1 - C_4 -alkyl;
 R_1 is C_1 - C_4 -alkyl or phenyl- C_1 - C_4 -alkyl; R_2 is hydrogen, halogen or C_1 - C_4 -alkoxy; R_3 is hydrogen or halogen;
 R_4 and R_6 , independently of one another, are hydrogen, halogen, or C_1 - C_4 -alkoxy; X_1 and X_3 are -O-;
 X_2 is C_4 - C_7 -alkylene; or a pharmaceutically acceptable salt thereof.

11. A compound according to claim 1 of formula IB




wherein R is hydrogen;

R_1 is C_1 - C_4 -alkyl; R_2 and R_3 , independently of one another, are hydrogen or halogen; X_1 and X_3 are -O-;
 X_2 is C_4 - C_7 -alkylene; or a pharmaceutically acceptable salt thereof.

12. A compound according to claim 5 which is 4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.


13. A compound according to claim 5 which is 4-[5-(5-fluoro-2-methoxyphenoxy)pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.

14. A compound according to claim 5 which is 4-[5-(4,5-difluoro-2-methoxyphenoxy)-pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.
15. A compound according to claim 5 which is 4-[5-(2,6-dimethoxyphenoxy)pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.
16. A compound according to claim 5 which is 4-[5-[2-(phenylmethoxy)phenoxy]pentoxy] benzenecarboximidamide or a pharmaceutically acceptable salt thereof.
17. Process for the manufacture of compounds of formula I defined in claim 1 and in which R is hydrogen and pharmaceutically acceptable salts thereof comprising
- a) converting in a compound of the formula IIa




or a salt thereof in which Z_1 is a radical which can be converted into the variable $-C(=NH)-NH-R$, Z_1 into the variable $-C(=NH)-NH-R$, or

- b) reacting a compound a compound of the formula IIIa



or a salt thereof with a compound of the formula IIIb



or a salt thereof in which Z_2 is a group of the formula $-X_1-X_2-Z_4$ and Z_3 is $-Z_5$, or Z_2 is $-Z_5$ and Z_3 is a group of the formula $Z_5-X_2-X_3$, wherein one of the radicals Z_4 and Z_5 is hydroxy or mercapto and the other is hydroxy, mercapto or reactive esterified hydroxy.

and, if desired, converting a compound of the formula I or a salt thereof obtainable according to the process or in another manner into another compound or a salt thereof according to the invention, converting a free compound of the formula I obtainable according to the process into a salt, converting a salt obtainable according to the process into the free compound of the formula I or into another salt, or resolving a mixture of isomers obtainable according to the process and isolating the desired compound.

18. A pharmaceutical preparation containing a compound of formula I according to claim 1 and customary carriers.
19. The use of compounds of formula I according to claim 1 or salts thereof and customary carriers for producing pharmaceutical preparations.
20. A compound of the formula I according to claim 1 or salts thereof for use in a method for the therapeutic

treatment of the human or animal body.

21. The compounds of the formula I according to claim 1 or salts thereof for use as antagonist of LTB_4 on human PMN's.


22. Use of a compound of formula I according to claim 1 for treating rheumatoid arthritis, inflammatory bowel disease, psoriasis, non-steroidal-antiinflammatory-drug-induced gastropathy, adult respiratory distress syndrome (ARDS), myocardial infarctions, allergic rhinitis, hemodialysis-induced neutropenia, and late phase asthma.

23. A pharmaceutical composition suitable for antagonizing LTB_4 receptors in mammals comprising an effective LTB_4 receptors antagonizing amount of a compound of claim 1 or a pharmaceutically acceptable salt thereof and a carrier

24. A method of the treatment of a condition or syndrome responsive to the selective antagonism of LTB_4 receptors which comprises administering to a mammal in need thereof an effective LTB_4 receptors antagonizing amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof of a said compound and a carrier.

Claims for the following Contracting States: ES, GR

1. A process for the manufacture of a compound of the formula I



wherein:

R is hydrogen or an acyl radical which is derived from an organic carbonic acid, an organic carboxylic acid, a sulfonic acid, or a carbamic acid;

R_1 is a substituent selected from an aliphatic hydrocarbon radical, an araliphatic hydrocarbon radical and a cycloaliphatic hydrocarbon radical;

X_1 and X_3 , independently of one another, are oxygen (-O-) or sulphur (-S-); and


X_2 is a divalent aliphatic hydrocarbon radical which may be interrupted by an aromatic radical;

wherein the phenyl rings of formula I may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid;

wherein aryl in the above definitions may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid; and


wherein a cycloaliphatic hydrocarbon radical may be substituted by an aliphatic radical; or a pharmaceutically acceptable salt thereof, comprising

a) converting in a compound of the formula IIa




10 or a salt thereof in which Z_1 is a radical which can be converted into the variable $-C(=NH)-NH-R$, Z_1 into the variable $-C(=NH)-NH-R$, or

b) reacting a compound of the formula IIIa



or a salt thereof with a compound of the formula IIIb



or a salt thereof in which Z_2 is a group of the formula $-X_1-X_2-Z_4$ and Z_3 is $-Z_5$, or Z_2 is $-Z_5$ and Z_3 is a group of the formula $Z_5-X_2-X_3$, wherein one of the radicals Z_4 and Z_5 is hydroxy or mercapto and the other is hydroxy, mercapto or reactive esterified hydroxy,

and, if desired, converting a compound of the formula I or a salt thereof obtainable according to the process or in another manner into another compound or a salt thereof according to the invention, converting a free compound of the formula I obtainable according to the process into a salt, converting a salt obtainable according to the process into the free compound of the formula I or into another salt, or resolving a mixture of isomers obtainable according to the process and isolating the desired compound.

2. A process according to claim 1 wherein there are produced compounds of formula I wherein R is hydrogen, (C_1-C_{12}) alkoxycarbonyl or (C_2-C_{12}) alkenyloxycarbonyl, each of which is unsubstituted or substituted by phenyl, naphthyl, indanyl or fluorenyl, or is (C_3-C_7) cycloalkoxycarbonyl being unsubstituted or mono- or polysubstituted by lower alkyl, or is lower alkanoyl or phenyl-lower alkanoyl, or is benzoyl, naphthoyl, indanoyl or fluorenyl, or is (C_1-C_7) alkanesulfonyl, phenyl- (C_1-C_7) alkanesulfonyl, (C_3-C_7) cydoalkanesulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by lower alkyl, phenyl-lower alkyl or phenyl;

R_1 is lower alkyl, lower alkenyl, lower alkynyl, phenyl-lower alkyl, phenyl-lower alkenyl, phenyl-lower alkynyl, (C_3-C_7) cycloalkyl, (C_3-C_7) cycloalkenyl, (C_3-C_7) cycloalkyl-lower alkyl or (C_3-C_7) -cycloalkenyl-lower alkyl;


X_1 and X_3 , independently of one another, are O or S;

X_2 is lower alkylene, lower alkylene-phenylene-lower alkylene or lower alkylene-naphthylene-lower alkylene;

wherein the phenyl rings of formula I may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, lower alkyl, lower alkenyl, lower alkynyl, hydroxy, lower alkoxy, lower alkenyloxy, phenyl-lower alkoxy, lower alkanoyloxy, lower alkenoyloxy and phenyl-lower alkanoyloxy;

wherein the aromatic radicals in the above definitions may be, independently of one another, substituted by one or more substituents selected from halogen, trifluoromethyl, lower alkyl, lower alkenyl, lower alkenyl, hydroxy, lower alkoxy, lower alkenyloxy, phenyl-lower alkoxy, cyclohexyl-lower alkoxy, lower alkanoyloxy, lower alkenyloxy and phenyl-lower alkanoyloxy; or a pharmaceutically acceptable salt thereof.


3. A process according to claim 1 wherein there are produced compounds of formula I wherein R is hydrogen, C₁-C₁₂-alkoxy-carbonyl, C₂-C₆-alkanoyl, phenyl-C₂-C₆-alkanoyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C₁-C₇-alkyl, or C₁-C₇-alkoxy, C₃-C₆-cycloalkylcarbonyl which is unsubstituted or substituted by C₁-C₇-alkyl, or is benzoyl, naphthoyl, indanoyl or fluorenyl, or is C₁-C₇-alkanesulfonyl, phenyl-C₁-C₇-alkanesulfonyl, C₃-C₇-cycloalkanesulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by C₁-C₇-alkyl, phenyl-C₁-C₇-alkyl or phenyl;
R₁ is C₁-C₇-alkyl, phenyl-C₁-C₇-alkyl or C₃-C₆-cycloalkyl;
X₁ and X₃ each are -O-, or furthermore are, independently of one another, -O- or -S-;
X₂ is C₂-C₇-alkylene or C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene;
wherein phenyl in the above definitions is unsubstituted or, furthermore, independently of one another, substituted by one or more substituents selected from halogen, trifluoromethyl, C₁-C₇-alkyl, and C₁-C₇-alkoxy; or a pharmaceutically acceptable salt thereof.
4. A process according to claim 1 wherein there are produced compounds of formula I wherein -O-R₁ is located in position 2 (ortho) of the corresponding phenyl ring with respect to -X₁-; and -C(=NH)-NH-R is located in position 4 (para) of the corresponding phenyl ring with respect to -X₃-; or a pharmaceutically acceptable salt thereof.
5. A process according to claim 1 wherein there are produced compounds of formula I of formula IA



- wherein R is hydrogen, C₁-C₁₂-alkoxycarbonyl, phenyl-C₁-C₄-alkoxycarbonyl, C₂-C₆-alkanoyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C₁-C₄-alkyl, or C₁-C₄-alkoxy, C₃-C₆-cycloalkylcarbonyl which is unsubstituted or substituted by C₁-C₄-alkyl;
R₁ is C₁-C₄-alkyl, phenyl-C₁-C₄-alkyl, C₃-C₆-cycloalkyl;
R₂, R₃, R₄ and R₅, independently of one another, are hydrogen, halogen, C₁-C₄-alkoxy, or C₁-C₄-alkyl;
X₁ and X₃ are -O-;
X₂ is C₄-C₇-alkylene or C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene; or a pharmaceutically acceptable salt thereof.
6. A process according to claim 1 wherein there are produced compounds of formula IA wherein R is hydrogen;
R₁ is C₁-C₄-alkyl, phenyl-C₁-C₄-alkyl, C₃-C₆-cycloalkyl;
R₂, R₃, R₄ and R₅, independently of one another, are hydrogen, halogen, C₁-C₄-alkoxy, or C₁-C₄-alkyl;
X₁ and X₃ are -O-;
X₂ is C₄-C₇-alkylene or C₂-C₄-alkylene-phenylene-C₂-C₄-alkylene; or a pharmaceutically acceptable salt thereof.
7. A process according to claim 1 wherein there are produced compounds of formula Ia wherein R is C₁-C₄-alkanesulfonyl, phenyl-C₁-C₄-alkanesulfonyl, C₃-C₇-cycloalkane-sulfonyl, or phenylsulfonyl, or is aminocarbonyl which is substituted by C₁-C₄-alkyl, phenyl-C₁-C₄-alkyl or phenyl;
R₁ is C₁-C₄-alkyl, phenyl-C₁-C₄-alkyl, C₃-C₆-cycloalkyl;
R₂, R₃, R₄ and R₅, independently of one another, are hydrogen, halogen, C₁-C₄-alkoxy or C₁-C₄-alkyl;
X₁ and X₃ are -O-;

X_2 is C_4 - C_7 -alkylene or C_2 - C_4 -alkylene-phenyl- C_2 - C_4 -alkylene; or a pharmaceutically acceptable salt thereof.

8. A process according to claim 1 wherein there are produced compounds of formula IA wherein
- 5 R is hydrogen;
 R_1 is C_1 - C_4 -alkyl or phenyl- C_1 - C_4 -alkyl;
 R_2 is hydrogen or C_1 - C_4 -alkoxy;
 R_3 is hydrogen or halogen;
 R_4 and R_6 , independently of one another are hydrogen, halogen or C_1 - C_4 -alkoxy;
 10 X_1 and X_3 are -O-;
 X_2 is C_4 - C_7 -alkylene; or a pharmaceutically acceptable salt thereof.
9. A process according to claim 1 wherein there are produced compounds of formula IA wherein R_4 and R_6 are hydrogen.
- 15 10. A process according to claim 1 wherein there are produced compounds of formula IA wherein R is C_1 - C_{12} -alkoxycarbonyl, phenyl- C_1 - C_4 -alkoxycarbonyl, C_2 - C_5 -alkanoyl, benzoyl which is unsubstituted or substituted by halogen, trifluoromethyl, C_1 - C_4 -alkyl, or C_1 - C_4 -alkoxy, C_3 - C_8 -cycloalkylcarbonyl which is unsubstituted or substituted by C_1 - C_4 -alkyl; R_1 is C_1 - C_4 -alkyl or phenyl- C_1 - C_4 -alkyl; R_2 is hydrogen, halogen or C_1 - C_4 -alkoxy; R_3 is hydrogen or halogen; R_4 and R_6 , independently of one another, are hydrogen, halogen, or C_1 - C_4 -alkoxy; X_1 and X_3 are -O-; X_2 is C_4 - C_7 -alkylene; or a pharmaceutically acceptable salt thereof.
- 20 11. A process according to claim 1 wherein there are produced of formula IB



- 35 wherein R is hydrogen;
 R_1 is C_1 - C_4 -alkyl; R_2 and R_3 , independently of one another, are hydrogen or halogen; X_1 and X_3 are -O-;
 X_2 is C_4 - C_7 -alkylene; or a pharmaceutically acceptable salt thereof.
- 40 12. A process according to claim 1 wherein there is produced 4-[5-(2-methoxyphenoxy)pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.
13. A process according to claim 1 wherein there is produced 4-[5-(5-fluoro-2-methoxyphenoxy)pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.
- 45 14. A process according to claim 1 wherein there is produced 4-[5-(4,5-difluoro-2-methoxyphenoxy)pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.
15. A process according to claim 1 wherein there is produced 4-[5-(2,6-dimethoxyphenoxy)pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.
- 50 16. A process according to claim 1 wherein there is produced 4-[5-[2-(phenylmethoxy)phenoxy]pentoxy]benzenecarboximidamide or a pharmaceutically acceptable salt thereof.



⑪ Publication number : **0 518 818 A3**

⑫ **EUROPEAN PATENT APPLICATION**

⑲ Application number : **92810422.3**

⑤① Int. Cl.⁵ : **C07C 257/18, A61K 31/155**

⑳ Date of filing : **02.06.92**

③① Priority : **11.06.91 US 714109**

⑦① Applicant : **CIBA-GEIGY AG**
Klybeckstrasse 141
CH-4002 Basel (CH)

④③ Date of publication of application :
16.12.92 Bulletin 92/51


⑦② Inventor : **Main, Alan Joseph**
60 Briar Lane
Basking Ridge, NJ 07920 (US)

⑧④ Designated Contracting States :
AT BE CH DE DK ES FR GB GR IT LI LU NL PT
SE

⑧⑧ Date of deferred publication of search report :
28.04.93 Bulletin 93/17

⑤④ Arylethers, their manufacture and use as medicament.

⑤⑦ The invention relates to the compounds of the formula



and pharmaceutically acceptable salts thereof, in which :

R is hydrogen or an acyl radical which is derived from an organic carbonic acid, an organic carboxylic acid, a sulfonic acid, or a carbamic acid ;

R₁ is a substituent selected from an aliphatic hydrocarbon radical, an araliphatic hydrocarbon radical and a cycloaliphatic hydrocarbon radical ;

X₁ and X₃, independently of one another, are oxygen (-O-) or sulphur (-S-) ; and

X₂ is a divalent aliphatic hydrocarbon radical which may be interrupted by an aromatic radical ;

wherein the phenyl rings of formula I may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid ;

wherein aryl in the above definitions may be, independently of one another, further substituted by one or more substituents selected from halogen, trifluoromethyl, an aliphatic hydrocarbon radical, hydroxy, and hydroxy which is etherified by an aliphatic, araliphatic or cycloaliphatic alcohol or which is esterified by an aliphatic or araliphatic carboxylic acid ; and wherein a cycloaliphatic hydrocarbon radical may be substituted by an aliphatic radical. The compounds are useful as selective LTB-4 receptor antagonists in the treatment of conditions or syndromes in mammals which are responsive to LTB-4 receptor antagonism.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application Number

EP 92 81 0422

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 5)
A	US-A-4 808 604 (A. BECK) * column 2, line 55 - line 63; claims *	1,18-24	C07C257/18 A61K31/155
A	EP-A-0 150 118 (MERCK FROSST CANADA INC) * page 4, line 26 - page 5, line 12; claims *	1,18-24	
A	EP-A-0 366 066 (R. R. TITWELL) * claims *	1	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 5)
			C07C A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims</p> <p>Claims searched completely:</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search THE HAGUE		Date of completion of the search 15 FEBRUARY 1993	Searcher SEUFERT G.H.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>A : member of the same patent family, corresponding document</p>			

EPF FORM 100 (12/92)



EP 92 81 0422 -C-

Remark : Although claim 24 is directed to a method of treatment of the human body (Art. 52(4) EPC) the search has been carried out and based on the alleged effects of the compounds/composition.